

PITNEY, HARDIN, KIPP & SZUCH, LLP

711 THIRD AVENUE, 20TH FLOOR

NEW YORK, NEW YORK 10017

PHONE : 212-687-6000

FACSIMILE : 212-682-3485

THE INFORMATION CONTAINED IN THIS FACSIMILE IS ATTORNEY PRIVILEGED AND CONFIDENTIAL INFORMATION INTENDED ONLY FOR THE USE OF THE INDIVIDUAL OR ENTITY NAMED BELOW. IF THE READER OF THIS FACSIMILE IS NOT THE INTENDED RECIPIENT OR THE EMPLOYEE OR AGENT RESPONSIBLE FOR DELIVERING IT TO THE INTENDED RECIPIENT, YOU ARE HEREBY NOTIFIED THAT ANY DISSEMINATION OR COPYING OF THIS FACSIMILE IS STRICTLY PROHIBITED. IF YOU HAVE RECEIVED THIS FACSIMILE IN ERROR, PLEASE NOTIFY US IMMEDIATELY BY TELEPHONE. THANK YOU.

x Please use the
attached page
rather than the
one previously sent
(typographical
error in
Figure 7)

FACSIMILE COVER PAGE

TO: YOUNG J. KIM

COMPANY: USPTO, Group Art Unit 1612

FACSIMILE NUMBER : 703-308-0294

DATE : January 11, 2001

FROM : Ronald R. Santucci

NO. OF PAGES : 2

Re: USSN 09/381,387

Our Docket No.: 2727-95

COMMENTS As requested accompanied herewith is a Brief Description of the Drawings for the above application. It can probably be inserted on page 3, after line 5, of the English translation unless you feel there is a more suitable place.

Regards,
Ronald Santucci

IF YOU HAVE ANY QUESTIONS REGARDING THIS TRANSMISSION, PLEASE CALL
(212)687-6000

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 is a graph of a FACS analysis showing the difference in fluorescence between two populations of agarose beads containing phosphorylated and non-phosphorylated sequences.

Fig. 2 is a graph of a FACS analysis showing the difference in fluorescence between two populations of agarose beads which were mixed and then treated with proteinkinase (PKA) and phosphorylated.

Fig. 3 is a FACS population distribution graph showing the two populations of beads, R2 and R3, which were sorted out by the FACS for further analysis.

Fig. 4 is a tabulation of the pool sequence analysis of the two group of beads, R2 and R3 which were sorted out of the population, as shown in Fig. 3.

Fig. 5 is a graph of the fluorescence of ovalbumin which was labeled with fluorescein/IDA/iron complex according to the invention and then separated by capillary electrophoresis.

Fig. 6 is a graph of the fluorescence of the control protein, glucose oxidase, which was similarly labeled as the ovalbumin in Fig. 5, and then separated by capillary electrophoresis.

Fig. 7 is a graph of a FACS analysis showing the difference in fluorescence between two populations of beads carrying an octapeptide, one population having a serine phosphate and the other an unmodified serine, following treatment with IDA and fluorescein as per the invention.

*** RX REPORT ***

RECEPTION OK

TX/RX NO	5016	
CONNECTION TEL		2126823485
SUBADDRESS		
CONNECTION ID		
ST. TIME	01/11 13:47	
USAGE T	00'49	
PGS.	2	
RESULT	OK	